


We Care more ...





REMEDIES PRIVATE LIMITED

29/2A, HAREKRISHTO SETT LANE, KOLKATA: 700 050 Ph.: 033 25589461/033 25322083 email: legendremedieskolkata@gmail.com website: www.lrpl.in

Head Office: UG-10, N. J. Complex Surat, Kadodara Road P.O.: Magob, State: Gujarat Pin: 395010, Ph.: 0261 2640581



Albendazole SAFEST DEWORMER

WORM INFESTATION IS THE MAIN REASON FOR PRODUCTION LOSS

BRAND IN MICRONIZED FORM

INDICATION:

For all types of Worm Infestation

DOSAGE: Bolus: 5 mg. to 7.5 mg. / kg. Body weight

Suspension:

Cattle: 1ml. for 5kg. Body wt.

Dog: 1ml./kg. Body wt. for 3 days

AVAILABLE:

Suspension:

30 ml./ 60 ml./ 90 ml./ 1 ltr.

Rolus :

150 mg./ 600 mg./ 1500 mg./ 3000mg.

DOSAGE:

INDICATION:

AVAILABLE: 1gm. -4 boli strip.

200 mg.-10 tab. strip.

ZANIDE

Oxyclozanide B.P. 200 mg. / 1000 mg. (VET)

THE MOST EFFECTIVE FLUKICIDE

For the treatment and control of fasciliosis.

Bolus: 1 gm.: 1 bolus for 66.6 kg. body wt.

Tablet: 200 mg.: 1 tab for 13.3 kg. body wt.

**Bolus & Tablet** 

Suspension

Levamisole Hydrochloride I.P.

A broad spectrum anthelmintic for treatment of parasitic infestations due to roundworms, lung worms, liver flukes and amphistomes.

INDICATION:

For all types of Worm Infestation

DOSAGE:

Sheep, Goats, Calves Cattle: 1 ml. / 2 Kg. body weight.

5 ml. / 10 Kg. body weight.

or as directed by Veterinarian

AVAILABLE:

Suspension: 90ml.

NENAZO

A broad spectrum Benzamidazole Anthelmintic

used against Gastro intestinal Parasitic including Giardia, Round Worm, Hook Worm, Whip Worm, Pin Worm and Taenia genus of Tape Worm. Nenazol can be administered to Sheep, Cattle, Dogs, Cats and Rabbits.

INDICATION:

A better solution to treat worm infestations.

DOSAGE:

Sheep, Goats, Calves Cattle: 5 mg. / Kg. body weight.

1.5 am. / 300 Kg. body weight. or as directed by the Veterinarian.

AVAILABLE:

1. Strip of 1 Bolus, 50 Strips in One Elegant Steel coloured Container.

2. 150 mg. Tablet (10 Tablets in one Blister).

SORBITOL BASED

Liquid Herbal Feed Supplement

THE HERBAL WONDER LIVER PROTECTANT

Fortified with

Kalmegh, Sarpankho, Bhrungraj, Kakamachi, Bhumyamlaki, Vrushya, Neem, Haridra, Kutaki, Sunthi, Darunisha, Kasani, Punarnava, Kumari, Vidanga, Jethimadh etc.

INDICATION:

Liver Cirrhosis, Enlargement of liver, Hepatitis, Anorexia, Jaundice and general debility.

DOSAGE:

For Calves: 10ml. daily for 5 days.

For Large Animal: 50ml. daily for 5 days. In acute condition, Please Prescribe the same dose for 10 days.

AVAILABLE:

100 ml., 200 ml., 500 ml., 1 Ltr. & 5 Ltr.

Injection

Liver Extract with B-Complex & Vitamin B12 Injection Fach ml. contains :

Thiamine Hydrochloride I.P. 25 mg. The Modern Liver

Total Vitamin B12

Chlorine Chloride

per ml.)

(Cyanocobalamin I.P. 30 mcg.

Riboflavin Sodium Phosphate I.P. 1.37 mg. | Extract Injection for Nicotinamide Pyridoxine Hydrochloride I.P.

I.P. 100 mg. <sub>1</sub> 5 mg. 32 mcg. 1

treating Liver Disorder INDICATION:

For the treatment of Liver Disorders.

& Liver Extract derived from 8 gm. of Fresh Liver having Cyanocobalamin activity equiv. to 2 mcg. Cyanocobalamin

Available: 15 mg, 30 ml./50 ml./100 ml.

Pnenol (as preservative) I.P. 0.5% w/v, i I.P.

Water for Injection

Q.S.

For Cattle, Buffaloes and Horses: (I / M) 5ml.-10 ml. daily for 3 or more days. For Cattle, Goats and Pigs: (I / M)

0.5ml,-2 ml, daily for 3 or more days. For Dogs & Cats: (I / M)

0.25ml.-2ml. daily for 3 or more days.

## LACTON Liquid

Herbal feed supplement for Animals for Improving Milk Yield

More MILK More PROFIT Milk enhancing formula

#### Fortified with

Jivanti, Satavari, Balant Soya, Amruta, Kamboji, Payaswini.

#### INDICATION:

Increases milk yield.

Increases the fat content in milk.

Protects animal from weakness in pregnancy. Improves Lactation.

Increases strength and immunity.

#### DOSAGE:

Large Animal: 100 ml. daily for 5-10 days. Small Animal: 50 ml. daily for 5-10 days.

#### AVAILABLE:

100ml., 500ml. &1 Ltr.

# MILCHEND Best Formulation for Quality Milk Yield

Powder

- INCREASE MILK BY SCIENTIFIC WAY
- THE ADVANTAGE OF CHELATED MINERALS FOR INFERTILITY

#### INDICATION:

Increases milk production. Increases SNF in milk.

#### DOSAGE:

Powder:

Large animal - 25 gms. Per day. Small animal - 5 gms. Per day.

#### AVAILABLE:

250 gm. powder, 1 kg. powder

A Super Premix of essential Vitamins, Amino Acids, Probiotics and Minerals for improving milk yield in Animals.

## **FIBROSIN**

Bolus

An Ayurvedic Veterinary Medicine to cure & control MASTITIS

#### Enriched with

Kanchnar gugal, Chitrak mula, Punarnavastaka, Trifala gugal & Apamarga

#### **INDICATION:**

Clinical & sub-clinical mastitis, As prophylactic, for effective control of mastitis, For cleansing of udder tissues, Co-therapy with antibiotics.

#### DOSAGE:

**TREATMENT**: At the first sight of Mastitis, 1st day to 3rd day 2 bolus twice stat. 4nd day to 6th day 1 bolus twice daily.

PREVENTION: One bolus twice daily for 5-6 days.

Herbal, Proven, Economic MASTITIS Therapy

AVAILABLE: 6 boli strip.

## **MOFULAG**

Liqui

### An Ayurvedic Veterinary Medicine Contains:

Vahaneri, Dikamali, Payaswini, Barash Karpur, Neem, Jethi Madh, Fatakadi & Nisha.

- STRESS due to transport, dietary, chemical exposure pollution and fatigue
- INAPPETANCE due to antibiotics, antibacterials anthelmentics & other chemical drugs
- NON SPECIFIC ANOREXIA

#### INDICATION:

Nonspecific Anorexia, Dyspepsia, Inappetance, Stress and supportive therapy in F.M.D. (Foot and Mouth Disease Virus)

#### DOSAGE

Large Animal: 50 ml. daily for 10 days. Small Animal: 10 ml. daily for 10 days.

#### AVAILABLE :

100 ml. / 500 ml. / 1 Litre

The Herbal Nutraceutical Immunity Improver And Stress Reducer

## **MOFULAG**

Granules

### An Ayurvedic Veterinary Medicine

- STRESS due to transport, dietary, chemical exposure pollution and fatigue
- INAPPETANCE due to antibiotics, antibacterials anthelmentics & other chemical drugs
- NON SPECIFIC ANOREXIA

### Contains

Vahaneri, Dikamali, Payaswini, Karpur, Neem Ext., Neem Patra, Godanti Bhasma.

#### INDICATION:

Nonspecific Anorexia, Dyspepsia, Inappetance, Stress and supportive therapy in F.M.D.\* \*(Food and Mouth Disease Virus)

#### DOSAGE

Large Animal: 30 gms. for 5 days with mollases. Small Animal: 10 gms. for 5 days with mollases.

#### AVAILABLE:

30 gm. in Tetra Pack, 100 gm. in Container TREAT THE SYMPTOMS OF F.M.D.

### CAAFOIL

Multi Action Skin Dressing Oil

The Best Herbal Dermatological Oil

#### Fortified with

Neem Oil, Karanj Oil, Nilagira Oil, Chalmogra Oil, Tarpin Oil, Panchagun Oil, Baras Karpur, Guda Pushpa Oil, Sharbana Oil.

### Ayurvedic Multi Action Skin Dressing Oil

### INDICATION:

Sarcoptic Mange, Psoroptic Mange, Demodectic Mange, Ringworm, Dermatomycosis, Eczema, Degnalla Surgical Wounds, Traumatic, Wounds, Lacerated Wounds, Broken Horn, Bite Wounds, Sinuses & Fistulae, Sores and Fissures, Yoke & Rope Gall, Infected Wounds, Septic Wounds, Gangrenous Wounds, Ulcerated Wounds, Foot Rot in Sheep, Foot, Lesions in F.M.D., Maggotted Wounds, Burn Wounds

#### DIRECTION:

Clean the affected area, and apply SAAF- OIL in sufficient quantity, once or twice daily until healing.

### AVAILABLE:

50ml., 100ml.

## **INSECTOR**

Composition:

Neem Tel 14.0% Karanja Tel 14.0% Nilgiri Tel 5.0%

0.3% Chaalmograa Terpentine Tel 7.0% 2.0% Barash Karpur

Gudapuspa 2.0% Sharbana 1.0% **HERBAL & SAFE** 

A Natural Herbal Remedy...

for healing of wounds and recovery from various skin infections.

> AVAILABLE: 100ml, Pack

All types of Wounds including Deep Seated Wounds. All types of Maggot Infestations Dermatitis. Surgical Incisions Foot Rot. Post

Operative dressings.



Norfloxacin Tinidazole 1800mg. (VET)

Mission to kill all causative agents for DIARRHOEA

The combination of Norfloxacin and Tinidazole has an extended range of Antimicrobial Spectrum and effective in mixed Gut Infections, Diarrhoea and Dysentery.

INDICATION:

Active against E.coli, Salmonella, Klebslella, Moraxella, Haemophilus, Anaerobic Bacteria, Clostridium Sp. and Giardiasis caused Diarrhoea and Dysentery.

DOSAGE:

2 Boli twice daily.

AVAILABLE:

4 Boli Strip

UMAR-NT

Tinidazole 600 mg. (VET)

INDICATION:

SPECIAL CARE FOR CALF DIARRHOEA

The combination of Norfloxacin and Tinidazole has an extended range of Antimicrobial Spectrum and effective in mixed Gut Infections, Diarrhoea and Dysentery.

INDICATION:

Treatment of Calf Scour, Diarrhoea and Dysentery of Amoebic Bacterial or Mixed Origins.

DOSAGE:

1 Tablet BD for 5 days.

AVAILABLE:

10 Tablets blister pack.

Trimethoprim Sulphamethoxazole I.P. 2000 mg. (VET)

Synergistic Combination of two powerful Antibacterials.

Hemorrhagic Septicemia (HS), Metritis, Acute and sub acute bacterial diarrhoea, Enteritis, Black Quarter (BQ), Secondary

Bacterial Infections, Upper Respiratory Tract Infections, Retained Placenta, Calf Scours.

- Can be safely used in pregnant animals.
- Doesn't interfere with calcium metabolism.

DOSAGE:

4 Boli at 12 hour interval for 3 - 5 days.

AVAILABLE:

4 Boli per strip.

## EMIDINE

Sulphadimidine I.P. 5 gm. (VET)

LEGEND'S PRIDE

Effective in treatment of infections caused by Gram (+ve) and Gram (-ve) bacteria and certain Protozoal diseases of livestock. Specially indicated for DIARRHOEA.

DOSAGE:

Large Animal: 100 mg. - 200 mg./ kg. body weight followed by half dose daily.

Small Animal: 1 gm/ 5 kg. body weight followed

by 0.5 gm/5 kg. daily 2 hourly.

AVAILABLE:

4 Boli per strip.

# MOXGEN-1.5 Bolus

NEW GENERATION AMOXYCILLIN

Amoxycillin Trihydrate I.P. equivalent to Amoxycillin 1500 mg. (VET)

**External Symptom of Respiratory Tract Infection** 

- Acid stable and well absorbed from gut.
- Higher peak serum level than Ampicillin after similar oral dose.
- Rapid excretion in the urine.
- Elimination half life 90 minutes in calf and dogs.

#### INDICATION:

Respiratory tract infections, Urinary Tract infection, Meningitis, Typhoid fever, Bacillary dysentery, Listeria infection, Infected bite wounds, Infected teeth, Bladder infections.

10 mg./kg. body weight twice daily or as directed by the Veterinarian.

AVAILABLE:

4 Boli in Strip

## GOFEED

Nutritional Supplement for Animal Feeding

#### Each Bolus Provides

Manganese Sulphate Dried Ferrous Sulphate Sodium Bi-carbonate Cobalt Chloride Zinc Sulphate Copper Sulphate Live yeast culture Liver Extract

#### INDICATION:

Improves apetite in anorexia caused due Calf/Sheep/Goat -1 bolus twice daily. to ruminal dysfunction. Improves 100 mg. ruminal movements and increases 500 mg. ruminal microflora. Can be used in 660 mg. correcting metabolic disoders such as 50 mg. ketosis and ruminal acidosis. Can be 100 mg. used in case of pubertal anoestrus and 250 mg. repeat breeding. Can be used as a 2 gm. supportive therapy during the usage of 5 gm. sulphonamides, antibiotics and Lactic Acid Bacilus 20,000,000 c.f.u. anthelmentics.

#### DOSAGE:

Cattle & Buffalowes - 2 boli twice daily. In case of anoestrous or repeat breeding 2 boli daily for 20 days.

### THE REAL APPETISER

#### AVAILABLE:

4 Boli blister pack.



Oxytetracycline 500 mg. (VET)

#### INDICATION:

Gram (+ve) and Gram (-ve) Bacteria, Spirochaetes, Rickettsia and certain Protozoa affecting livestock.

#### DOSAGE:

1 Tablet for 50-80 Kg. body weight.

#### AVAILABLE:

4 Tablet Strip

# L-EN-RO Injection

Enrofloxacin 10% w/v (VET)

4th Generation Floroquinolone for Modern Combat

L-EN-RO has a very broad antibacterial spectrum with very low MIC values, covering Gram (+ve), Gram (-ve) and other bacterial infections like Mycoplasma, Chlamydia etc.

#### INDICATION:

Bronchopneumonia, Contagious Bovine and Caprine Pleuropneumonia (Mycoplasma sp.), Dermatitis, Mastitis, Metritis, Pyometra, Wooden Tongue, Lumpy Jaw, Brucellosis, Abcess and any non specific or mixed infections where clinical picture is not clear.

#### DOSAGE:

Large Animal: 15 ml. daily for 3-5 days. Small Animal: 2 ml. - 3 ml. daily for 3-5 days.

#### AVAILABLE:

15 ml., 50 ml. & 100 ml. vial.

## LOFAC-MP

The FASTEST & SAFEST Fever and Pain Reliever.

A Potent NSAID Formulation.

Each ml. contains: Paracetamol 150 mg. 5 mg. (VET) Meloxicam

#### INDICATION:

Control of pain, fever and inflammation associated with Osteoarthritis, Myocitis, Mastitis, Metritis, Pneumonia, Dystokia, Fractures and Surgery.

#### DOSAGE:

Cattle, Horse, Camel, Elephant and other: 15ml.-30 ml. per 300 kg. Body Weight depending upon the severity of cases.

Calves, Sheep and Goat: 2 ml. per 10 kg. body weight

#### AVAILABLE:

Administration By I / M route only. 30 ml. vial. / 100 ml. vial.

### LOFAC-MP

Meloxicam I.P. 100 ma. Paracetamol I.P. 1500 mg. (VET)

> The Fastest & Safest Fever and Pain Reliever

Pneumonia, Osteoarthritis, Broncho Pneumonia, Fractures & Surgeries, Sprains & Strains, Prolapse of Uterus, Metritis, Dystokia, Arthritis, Myocitis, Mastitis, Joint Pain, Laminitis.

#### DOSAGE:

Large Animal: 2 Bolus / 400 Kg. Body Weight once a day, or as directed by the Veterinarian. Small Animal: 1/2 Bolus daily or as directed by the Veterinarian.

#### AVAILABLE :

4 Boli in Strips.

# LOFAC-M Bolus

Meloxicam 100 mg. (VET)

### THE ASSURED PAIN MANAGER

- Superior GI tolerability.
- Environmental Friendly.
- · No renal Problem.

#### INDICATION:

LOFAC-M relieves painful inflammatory conditions like muscular pain, joint pain, arthritis, sprains and

LOFAC-M relieves acute pain associated with dystokia, colic, mastitis, metritis, prolapse and fever.

LOFAC-M relieves painful inflammatory conditions in post parturition, post operative and post traumatic.

### DOSAGE:

Large Animals: 1 - 2 boli daily. Small Animals: 1/2 bolus daily.

#### AVAILABLE:

4 Boli blister pack.

## BLOTEND

Herbal wonder in the treatment of Tympanites, Flatulence and as a supportive therapy in constipation and indigestion

Active enzymes and alkaloids present in ingredients

Suwa, Ajwain, Jirak, Rason, Haritaki & Bibhitaki, Amalaki, Hing ras, Sunthi, Maricha and Pipper, Neem, Dikamali, Kalmegh.

#### INDICATION:

Bloat, Flatulence, Constipation and Indigestion.

Large Animals: 50 ml. twice daily. Small Animals: 25 ml. twice daily.

#### AVAILABLE:

100ml. Pet Bottle

# EMILYTE O.R.S. Powder

Each 5 gm. contains Sodium Chloride 40 mg. 250 mg. Potassium Chloride 150 mg. Sodium Bicarbonate Sodium Acid Phosphate 40 mg. Sodium Citrate 325 mg. Calcium Lactate 85 mg. 50 mg. Magnessium Sulphate Ascorbic Acid (coated) 65 mg. Dextrose a.s.

Revitalise your animal with O.R.S. Powder (VET) As a supportive therapy in E. Coli & Salmonella Infections in Poultry

INDICATION:

To provide instant energy. To rehydrate during dehydration and heat stress in summer. To combat against fluid and electrolytic loss due to diarrhoea and dysentery as a result of sulmonellosis, colibacilosis, John's disease, Coccidiosis. After I.V. dextrose inj.

#### DOSAGE:

Large Animal: 50 gm. twice daily for 4 days. Small Animal: 25 gm. twice daily for 4 days.

#### AVAILABLE:

25 ams / 200 ams Pouch

## **EDEXIN**

Cephalexin IP.

#### ANTI BACTERIAL UTERUS CLEANSER

- Broad spectrum and bactericidal.
- Achives peak serum level fast.
- Works effectively in the presence of pus, necrotic tissues, enzymes, bacteria or change of pH in infective uterus.

#### INDICATION:

Infections in Uterus causing ENDOMETRITIS, CERVICITIS, PYOMETRA.

#### DOSAGE:

1 Sachet to be dissolved in 50 ml. or regd, quantity of distilled water and to be given through intra-uterine route.

#### AVAILABLE:

20 gm. sachet.



Chick Salmonella. **E.coli and Bacillary** Diarrhoea, Wing Rot & Coryza

## OVATONE

The Herbal Ovary Toner Each capsule contains:

Mrugakshi Ext. A.B. 200 mg. Sunthi A.B. 60 mg. Dharampattan A.B. 60 mg. A.B. 60 mg. Vaidehi A.B. 10 mg. I Hing Ras

INDICATION:

Infertility due to ovarian dysfunction such as anoestrous, silent heat, delayed maturity in heifers. To reduce unproductive inter calving period.

### **Excipients** DOSAGE:

Large Animals: 6 caps /day. Small Animals: 2 caps /day.

Orally once daily for two consecutive days. In case of no oestrus or weak oestrus is observed, repeat the same dose after 10 days following the first treatment.

Q.S.

#### AVAILABLE:

6 Capsules in a strip

# VITAMIN-A

Each 2 ml. contains :

Vitamin A (Palmitate) 600000 I.U.

#### Prevention and Treatment in Hypovitaminosis A

Promotes growth. Improves appetite. Assists in reproduction and Lactation. Helps in normal vision. Treats infertility. It Stimulates germinal epithelium and helps in ovulation.

#### INDICATION:

Night Blindness, Stratified Keratinisation of epithelial tissue, Emaciation, Lacrimation, Xeropthalmia, Corneal cloudiness, Decline in male sexuality, Shortened gestation period, Retained placenta, Reduced milk yield & Infertility.

DOSAGE: Through I.M. route

Infertility (Male/Female): 6 ml. 1st and 3rd day

Large Animal: 12 ml. - 18 ml. Per Week Small Animal: 3 ml. - 6 ml. per week Canine, Feline: 2 ml. - 6 ml. per week AVAILABLE: 2 ml. x 6 ampoules in a box.

# MILCHEND Bolus

### THE ADVANTAGE OF CHELATED MINERALS FOR INFERTILITY

A Super Premix of Essential Vitamins, Amino Acids, Probiotics and Chelated Minerals for Improving Health & Fertility in Animals.

#### INDICATION:

Treats infertility. Helps in reproduction. Overcomes mineral deficiency. Maintains balance of nutrients during Pregnancy.

#### DOSAGE:

Prevention: 1-2 Bolus daily. Treatment: 2-4 Bolus daily.

#### AVAILABLE:

10 x 4 Boli.